

4. Conclusions

In this work, a highly efficient procedure for the synthesis of 2-amino-4H-chromenes in the presence of TiO₂/NCP as a heterogeneous and reusable nanocatalyst was performed under solvent-free condition. This procedure offers several merits including high yields of products and mild reaction conditions. Moreover, the obtained results demonstrated good reusability of the catalyst.

Acknowledgements

Partial financial support from the Research Council of Hakim Sabzevari University is greatly appreciated.

References

- [1] L. Edjlali, R.H. Khanamiri, *Monatsh. Chem.* (2016) 1-5.
- [2] S.M. George, *Chem. Rev.* 95 (1995) 475-476.
- [3] M.G. Dekamin, M. Eslami, A. Maleki, *Tetrahedron* 69 (2013) 1074-1085.
- [4] S. Gao, C.H. Tsai, C. Tseng, C.-F. Yao, *Tetrahedron* 64 (2008) 9143-9149.
- [5] I. Devi, P.J. Bhuyan, *Tetrahedron Lett.* 45 (2004) 8625-8627.
- [6] A.M. Shestopalov, Y.M. Emeliyanova, A.A. Shestopalov, L.A. Rodinovskaya, Z.I. Niazimbetova, D.H. Evans, *Org. Lett.* 4 (2002) 423-425.
- [7] M.C. Bagley, J.W. Dale, J. Bower, *Chem. Commun.* (2002) 1682-1683.
- [8] F. Javadi, R. Tayebee, *Microporous Mesoporous Mater.* 231 (2016) 100-109.
- [9] Y. Sarrafi, E. Mehrasbi, A. Vahid, M. Tajbakhsh, *Chin. J. Catal.* 33 (2012) 1486-1494.
- [10] V.M. Joshi, R.L. Magar, P.B. Throat, S.U. Tekale, B.R. Patil, M.P. Kale, R.P. Pawar, *Chin. Chem. Lett.* 25 (2014) 455-458.
- [11] S.S. Katkar, M.K. Lande, B.R. Arbad, S.T. Gaikwad, *Chin. J. Chem.* 29 (2011) 199-202.
- [12] A.S. Waghmare, S.S. Pandit, *Iran. Chem. Commun.* 3 (2015) 291-301.
- [13] S. Khaksar, A. Rouhollahpour, M. Saeed, *J. Fluorine Chem.* 141 (2012) 11.
- [14] J.-C. Xu, W.-M. Li, H. Zheng, Y.-F. Lai, P.-F. Zhang, *Tetrahedron* 67 (2011) 9582-9587.
- [15] R.J. Kalbasi, N. Mosaddegh, *Catal. Commun.* 12 (2011) 1231-1237.
- [16] M.N. Elinson, A.I. Ilovaisky, V.M. Merkulova, P.A. Belyakov, A.O. Chizhov, G.I. Nikishin, *Tetrahedron* 66 (2010) 4043-4048.
- [17] S. Tu, H. Jiang, Q. Zhuang, C. Miao, D. Shi, X. Wang, Y. Gao, *Chin. J. Org. Chem.* 23 (2003) 488-490.
- [18] T.-S. Jin, A.-Q. Wang, F. Shi, L.-S. Han, L.-B. Liu, T.-S. Li, *Arkivoc* 14 (2006) 78-86.
- [19] W.-B. Sun, P. Zhang, J. Fan, S.-H. Chen, Z.-H. Zhang, *Synth. Commun.* 40 (2010) 587-594.
- [20] M.R. Naimi-Jamal, S. Mashkouri, A. Sharifi, *Mol. Diversity* 14 (2010) 473-477.
- [21] S. Balalaie, M. Bararjanian, M.S. Ahmadi, S. Hekmat, P. Salehi, *Synth. Commun.* 37 (2007) 1097-1108.
- [22] M. Seifi, H. Sheibani, *Catal. Lett.* 126 (2008) 275-279.
- [23] J.M. Khurana, S. Kumar, *Tetrahedron Lett.* 50 (2009) 4125-4127.
- [24] J.T. Li, W.Z. Xu, L.C. Yang, T.S. Li, *Synth. Commun.* 34 (2004) 4565-4571.
- [25] R. Hekmatshoar, S. Majedi, K. Bakhtiari, *Catal. Commun.* 9 (2008) 307-310.
- [26] M.M. Heravi, B.A. Jani, F. Derikvand, F.F. Bamoharram, H.A. Oskooie, *Catal. Commun.* 10 (2008) 272-275.
- [27] B.S. Kumar, N. Srinivasulu, R. Udipi, B. Rajitha, Y.T. Reddy, P.N. Reddy, P. Kumar, *J. Heterocycl. Chem.* 43 (2006) 1691-1693.
- [28] G. Sabitha, K. Arundhathi, K. Sudhakar, B. Sastry, J. Yadav, *Synth. Commun.* 39 (2009) 433-442.
- [29] D. Fang, H.B. Zhang, Z.L. Liu, *J. Heterocycl. Chem.* 47 (2010) 63-67.
- [30] S. Makone, S. Mahurkar, *Green Sustainable Chem.* 3 (2013) 27-32.
- [31] M. Bahrami, A. Nezamzadeh-Ejehieh, *Mater. Sci. Semicond. Process.* 30 (2015) 275-284.
- [32] N. Rajic, D. Stojakovic, N. Daneu, A. Recnik, *J. Phys. Chem. Solids* 72 (2011) 800-803.
- [33] A. Godelitsas, T. Armbruster, *Microporous Mesoporous Mater.* 61 (2003) 3-24.
- [34] M. Dalconi, A. Alberti, G. Cruciani, P. Ciambelli, E. Fonda, *Microporous Mesoporous Mater.* 62 (2003) 191-200.
- [35] R. Tayebee, F. Javadi, G. Argi, *J. Mol. Catal. A: Chem.* 368 (2013) 16-23.
- [36] D.W. Breck, *Zeolite Molecular Sieves*, John Wiley & Sons, New York, 1974.
- [37] D. Kallo, J. Papp, J. Valyon, *Zeolites* 2 (1982) 13-16.
- [38] M.M. Treacy, J.B. Higgins, *Collection of Simulated XRD Powder Patterns for Zeolites*, Elsevier, 2007.
- [39] K. Thamaphat, P. Limsuwan, B. Ngotawornchai, *Kasetsart J. Nat. Sci.* 42 (2008) 357-361.
- [40] Y. Ren, F. Zhang, W. Hua, Y. Yue, Z. Gao, *Catal. Today* 148 (2009) 316-322.
- [41] F. Iacomi, *Surf. Sci.* 532 (2003) 816-821.
- [42] A. Pourtaheri, A. Nezamzadeh-Ejehieh, *Spectrochim. Acta Mol. Biomol. Spectrosc.* 137 (2015) 338-344.
- [43] J. Farzad, R. Tayebee, B. Bahramian, *Appl Organometal Chem.* (2017) doi: 10.1002/aoc.3779.
- [44] A. Nezamzadeh-Ejehieh, F. Khodabakhshi-Chermahini, *Ind. Eng. Chem. Res.* 20 (2014) 695-704.
- [45] S. Nemouchi, R. Boulcina, B. Carboni, A. Debache, *C.R. Chim.* 15 (2012) 394-397.
- [46] L.-M. Wang, J.-H. Shao, H. Tian, Y.-H. Wang, B. Liu, *J. Fluorine Chem.* 127 (2006) 97-100.
- [47] D. Shi, J. Mou, Q. Zhuang, X. Wang, *J. Chem. Res.* 2004 (2004) 821-823.
- [48] K. Niknam, N. Borazjani, R. Rashidian, A. Jamali, *Chin. J. Catal.* 34 (2013) 2245-2254.
- [49] M. Makvandi, F.A. Dil, A. Malekzadeh, M. Baghernejad, K. Niknam, *Iran. J. Catal.* 3 (2013) 221-228.